

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number
WO 03/086387 A1

(51) International Patent Classification⁷: **A61K 31/366**,
9/20

(21) International Application Number: PCT/CA03/00479

(22) International Filing Date: 3 April 2003 (03.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2,379,887 9 April 2002 (09.04.2002) CA

DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant and

(72) Inventor: **SHERMAN, Bernard, Charles** [CA/CA]; 50
Old Colony Road, Toronto, Ontario M2L 2K1 (CA).

Published:

— with international search report

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CH, CN, CO, CR, CU, CZ,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/086387 A1

(54) Title: STABLE TABLETS COMPRISING SIMVASTATIN

(57) Abstract: A tablet comprising simvastatin and excipients, wherein the content of lactose, if any, is less than 75 percent of the total excipients by weight, and wherein the content of cellulose is more than 20 percent of the total excipients by weight.

STABLE TABLETS COMPRISING SIMVASTATIN**BACKGROUND OF THE INVENTION**

5 Simvastatin is a lipid-lowering agent that is produced synthetically from lovastatin. Simvastatin is disclosed and claimed in U.S. Patent No. 4444784. Tablets comprising simvastatin as the active ingredient are sold by Merck & Co., Inc. in the United States and elsewhere under the tradename Zocor™ in strengths of 5 mg, 10 mg, 20 mg, 40 mg and 80 mg.

10

With respect to pharmaceutical formulations (i.e. dosage forms) comprising simvastatin, the only information disclosed in U.S. Patent No. 4444784 is a statement that typical formulations for filling hard gelatin capsules comprise the active drug and finely divided lactose.

15

20

Zocor™ tablets are film-coated tablets, by which is meant that they consist of core tablets surrounded by a water-soluble film coating. The labelling for Zocor™ tablets indicate that the excipients (i.e. inactive ingredients) used in the core tablets are lactose, cellulose, starch, magnesium stearate, ascorbic acid, citric acid and butylated hydroxyanisole (also known as BHA).

25

Lactose, cellulose, starch, and magnesium stearate are all very commonly used as excipients in tablets. In particular, lactose and cellulose are very commonly used as fillers and binders. When lactose is used as an excipient, it is typically used in the largest quantity and typically constitutes more than 75 percent of the total excipients by weight. Starch is very commonly used as a filler and disintegrant, and magnesium stearate is very commonly used as a lubricant, to avoid sticking and binding in the tableting process.

30

5 The inclusion of BHA, ascorbic acid, and citric acid as excipients in the core tablets is unusual, and the inclusion of these excipients indicates that Merck & Co. Inc. found that it was necessary to include these excipients to achieve satisfactory stability of the simvastatin in the tablet. Simvastatin is prone to degradation due to oxidation of the diene and oxidation of the hydroxyl group in the simvastatin molecule. BHA and ascorbic acid are apparently included in the tablets as antioxidants. Citric acid is apparently added because it has chelation properties with metal ions, which, in the absence of the citric acid, could catalyze the oxidation process.

15 The composition of the Zocor™ core tablets is thus relatively complex in terms of the number of excipients used. Also, the use of BHA as an antioxidant usually requires the use of a solvent. That is to say, BHA is dissolved in a solvent, the solution is used to granulate the simvastatin, optionally after mixing with one or more excipients, and the wet mass is then dried to evaporate the solvent. The process of manufacture of the tablets is thus much more complex and expensive than a simple dry-mix process, by which is meant a process in which all of the ingredients are mixed together in dry form, and the mixture is compressed into tablets, without adding a solvent and then drying to evaporate the solvent. It is clearly preferable to avoid the use of solvents, if possible, in order to simplify the process of manufacture.

25 Simvastatin is also a compound for which it is difficult to produce a tablet formulation which exhibits rapid absorption after ingestion. It is necessary that any tablet formulation that is developed as an alternative to Zocor™ exhibit a rate of absorption on oral administration that is equivalent to that of Zocor™.

30

In light of this prior art, the objective of the present invention is to enable the manufacture of simvastatin tablets so as to achieve at least one or more, if not all, of the following:

5

1. Improved stability relative to tablets that use lactose as the predominant excipient.
2. Rate and extent of absorption equivalent to Zocor™ upon oral administration.
3. Elimination of the need to include citric acid as an excipient.
4. Elimination of the need to include ascorbic acid as an excipient.
5. Elimination of the need to include BHA as an excipient.
6. Production by a dry-mix method; i.e. without the need to granulate with a solvent and then dry to evaporate the solvent.

10

15

20

BRIEF SUMMARY OF THE INVENTION

It has been found that rate of degradation of simvastatin is significantly affected by the excipients with which it is mixed.

25

More particularly, it has been found that the stability of simvastatin in tablets is significantly improved by reducing or eliminating the lactose content, and by using cellulose, as a major excipient. The tablets of the present invention thus comprise simvastatin and excipients, wherein the content of lactose, if any, is less than 75 percent of the total excipients by weight, and wherein the content of cellulose is more than 20 percent of the total excipients by weight.

30

DETAILED DESCRIPTION OF THE INVENTION

5 Tablets comprising simvastatin will generally be made by mixing simvastatin with excipients (inactive ingredients) and compressing the mixture into tablets on a tablet press.

10 Among ingredients most commonly used as fillers and binders in pharmaceutical tablets are lactose (which may be either anhydrous lactose or lactose monohydrate) and cellulose. They are considered to be binders as well as fillers, because they usually enable compression into hard tablets, if they are the predominant ingredients.

15 Because lactose is the predominant excipient used in Zocor™ tablets, which presumably were carefully developed to Merck & Co. Inc. for maximum stability, it has been surprising to discover that stability is better with cellulose than with lactose as principal excipient.

20 Tablets of the present invention will have a lactose content that is less than 75 percent, is preferably less than 60 percent, and is more preferably less than 40 percent of the total excipient content by weight. Most preferably the tablets will be lactose free.

25 The tablets will comprise cellulose (which may be either microcrystalline cellulose or powdered cellulose) as a filler and binder. The amount of cellulose will exceed 20 percent, will preferably exceed 40 percent, and will more preferably exceed 60 percent of the total excipients by weight.

30 Cellulose is often considered to be a disintegrant in addition to being a filler and binder, because, like other disintegrants, it absorbs water and swells, thus aiding in the disintegration of tablets containing cellulose when they

are added to an aqueous medium. Nevertheless, tablets of the present invention will preferably also comprise a disintegrant other than cellulose.

5 Commonly used disintegrants include starch (which may be pregelatinized), croscarmellose sodium, carmellose calcium, sodium starch glycolate, and crospovidone. It has been further found that stability of the simvastatin tablets is better when starch, sodium starch glycolate, or crospovidone is
10 used as disintegrant, than when either croscarmellose sodium or carmellose calcium is used. Accordingly, the tablets of the present invention will preferably comprise starch, sodium starch glycolate or crospovidone, and will preferably not comprise any croscarmellose sodium or carmellose calcium.

15 The presence of an adequate amount of a disintegrant other than cellulose is necessary to ensure that the tablets are bioequivalent to Zocor™; that is to say that the rate of absorption of the simvastatin is equivalent to that of Zocor™ upon ingestion.

20 The United States Pharmacopoeia, 25th edition, includes a dissolution test for simvastatin tablets. The test is done in apparatus 2 at 50 rpm in 900 mL of pH 7.0 buffer solution containing 0.5 percent dodecyl sodium sulfate in 0.01M sodium phosphate. The specification requires dissolution of not less
25 than 75 percent in 30 minutes.

While tablets which comprise cellulose as the predominant excipient and which do not also comprise another disintegrant may exhibit rapid disintegration in water and may even pass the United States Pharmacopoeia dissolution test, such tablets will nevertheless be unlikely to be bioequivalent
30 to Zocor™ tablets, as a result of exhibiting slower dissolution than Zocor™ in gastric fluid, which has a lower pH than used in the United States Pharmacopoeia test.

Where starch is used as the disintegrant, the quantity will preferably exceed 12 percent and more preferably exceed 20 percent of the total of excipients by weight. Where the disintegrant is selected from sodium starch glycolate, crospovidone, croscarmellose sodium or carmellose calcium (of which the first two of these are preferable as aforesaid for reasons of stability), the amount will preferably exceed 1 percent, more preferably exceed 2 percent, and even more preferably exceed 3 percent of the total of excipients by weight.

The total of excipients selected from the group consisting of cellulose, starch, sodium starch glycolate will preferably exceed 65 percent, will more preferably exceed 80 percent, will even more preferably exceed 90 percent, and will most preferably exceed 95 percent of the total of excipients by weight.

It has been found that the stability of simvastatin in tablets is also significantly affected by the choice of lubricant. Stability is improved when magnesium stearate, which is by far the most commonly used lubricant, is replaced by zinc stearate, or sodium stearyl fumarate. Thus tablets of this present invention will preferably be free of magnesium stearate, and will preferably comprise zinc stearate or sodium stearyl fumarate.

By selecting excipients in accordance with the teaching of this disclosure, it is possible to significantly improve the stability of simvastatin so as to reduce or eliminate the need for the stabilizers that are included in Zocor™ tablets.

Tablets of the present invention will thus optionally and preferably be free of citric acid. The tablets will also optionally and preferably be free of ascorbic acid. The tablets will preferably be free of both citric and ascorbic acid. The tablets will also optionally and preferably be free of BHA. The tablets will preferably be free of all three of citric acid, ascorbic acid and BHA.

As aforesaid, the inclusion of BHA in Zocor™ tablets requires the use of a wet-granulation process in which the BHA is dissolved in solvent, the solution is used to wet granulate the simvastatin (after it is mixed with excipients), and the wet mass is then dried to evaporate the solvent.

The tablets of the present invention will preferably be made by a dry-mix process; that is to say, a process in which all of the ingredients are mixed together in dry form, without any step of adding solvent and then drying to evaporate the solvent.

The invention will be better understood from the following examples, which are meant to be illustrative, and not limiting of the scope of the invention.

EXAMPLES 1 TO 7

Example No.	1	2	3	4	5	6	7
Simvastatin	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Anhydrous Lactose	39.6	25.0	25.0	25.0	25.0	25.0	25.0
Microcrystalline Cellulose	0	14.6	0	0	0	0	0
Starch, Pregelatinized	0	0	14.6	0	0	0	0
Croscarmellose Sodium	0	0	0	14.6	0	0	0
Carmellose Calcium	0	0	0	0	14.6	0	0
Sodium Starch Glycolate	0	0	0	0	0	14.6	0
Crospovidone	0	0	0	0	0	0	14.6
Magnesium Stearate	0.4	0.4	0.4	0.4	0.4	0.4	0.4
	45.0	45.0	45.0	45.0	45.0	45.0	45.0

For each of examples 1 to 7, the ingredients were mixed in the proportions shown, and the powder mixture was then compressed into tablets of unit weight 45 mg, so that each tablet comprised about 5 mg of simvastatin. In the tablets of example 1, the only excipients are lactose as filler-binder, and magnesium stearate as lubricant. In each of examples 2 to 7, 14.6 mg of the

lactose is replaced by 14.6 mg of one of cellulose, starch, croscarmellose sodium, carmellose calcium, sodium starch glycolate and crospovidone.

5 Tablets of each of examples 1 to 7 were stored at 60°C for four weeks and then tested for the amount of simvastatin-oxolactone, which is the product of oxidation of the hydroxyl group in the simvastatin molecule. The testing was done by an HPLC (high performance liquid chromatographic) method, with results as follows, as a percentage of the simvastatin content:

10

Example No.	1	2	3	4	5	6	7
% Oxolactone	.108%	.104%	.154%	.169%	.187%	.142%	.118%

15

The initial level of oxolactone in the simvastatin used to make the tablets was about 0.04%.

Comparing the results for examples 1 and 2 indicates that replacing part of the lactose by cellulose reduced the rate of oxidation.

20

Comparing the results of examples 3 to 7 with the result of example 1 indicates that the addition of any of the five disintegrants increases the rate of oxidation (at least when lactose is still the primary excipient), but that the rate of oxidation is lower when the disintegrant is crospovidone, sodium starch glycolate, or starch, than when it is croscarmellose sodium or

25

carmellose calcium.

30

EXAMPLES 8 TO 13

Example No.:	8	9	10	11	12	13
5 Simvastatin	5.0	5.0	5.0	5.0	5.0	5.0
Microcrystalline Cellulose	39.9	25.3	25.3	25.3	25.3	25.3
Starch, Pregelatinized	0	14.6	0	0	0	0
Croscarmellose Sodium	0	0	14.6	0	0	0
Carmellose Calcium	0	0	0	14.6	0	0
10 Sodium Starch Glycolate	0	0	0	0	14.6	0
Crospovidone	0	0	0	0	0	14.6
Magnesium Stearate	0.1	0.1	0.1	0.1	0.1	0.1
	45.0	45.0	45.0	45.0	45.0	45.0

Examples 8 to 13 are repeats of examples 2 to 7, with all of the lactose replaced by cellulose, and the amount of magnesium stearate per tablet reduced from 0.4 mg to 0.1 mg. The tablets were made by the same process as used in examples 1 to 7.

Sample tablets were also stored at 60°C for 4 weeks and tested by the same method, with results as follows:

Example No.	8	9	10	11	12	13
% Oxolactone	.098%	.113%	.129%	.143%	.101%	.095%

Comparing the result for example 8 to the results for examples 1 and 2 confirms that the rate of oxidation is further reduced by replacing the balance of the lactose with more cellulose. Comparing the results of examples 9 to 13 again confirms that the rate of oxidation is lower when the disintegrant is one of crospovidone, sodium starch glycolate or starch, than when it is croscarmellose sodium or carmellose calcium.

EXAMPLES 14 AND 15

	<u>Example No.</u>	<u>14</u>	<u>15</u>
5	Simvastatin	5.0	5.0
	Crospovidone	34.9	34.9
	Magnesium Stearate	0.1	0
	Zinc Stearate	<u>0</u>	<u>0.1</u>
		40.0	40.0

10 Tablets of examples 14 and 15 were made by the same process as examples 1 to 13, except that the tablets were made at a unit weight of 40 mg instead of 45 mg. The purpose of examples 14 and 15 was to compare the effect on oxidation rate when magnesium stearate is replaced by zinc
15 stearate as lubricant.

Again tablets were stored at 60°C, and after 4 weeks samples were tested with results as follows:

	<u>Example No.</u>	<u>14</u>	<u>15</u>
20	% Oxolactone	0.193%	0.146%

These results confirm that the oxidation rate is reduced by eliminating magnesium stearate as lubricant, and replacing it by zinc stearate.

25 Comparing the results of examples 14 and 13, also shows that the oxidation rate is lower when the excipient content is mostly cellulose than when it is mostly crospovidone.

30

EXAMPLES 16 TO 21

5	<u>Example No.</u>	<u>16</u>	<u>17</u>	<u>18</u>	<u>19</u>	<u>20</u>	<u>21</u>
	Simvastatin	5.0	5.0	5.0	5.0	5.0	5.0
	Microcrystalline Cellulose	39.9	30.0	39.9	30.0	40.0	30.0
	Crospovidone	0	9.9	0	9.9	0	10.0
	Zinc Stearate	0.1	0.1	0	0	0	0
10	Sodium Stearyl Fumarate	0	0	0.1	0.1	0	0
		45.0	45.0	45.0	45.0	45.0	45.0

Tablets of examples 16 to 21 were made by the same process as examples 1 to 13. The purpose of these examples was to compare the stability of tablets with zinc stearate as lubricant, sodium stearyl fumarate as lubricant, and no lubricant at all, all both with and without crospovidone as disintegrant.

Tablets for each of examples 16 to 21 were stored at 60°C for 2 weeks, and then tested with results as follows:

20

<u>Example No.</u>	<u>16</u>	<u>17</u>	<u>18</u>	<u>19</u>	<u>20</u>	<u>21</u>
% Oxolactone	.055%	.055%	.046%	.051%	.046%	.041%

These results indicate good stability, regardless of whether the tablets contain sodium stearyl fumarate or zinc stearate as lubricant, or no lubricant at all.

30

CLAIMS

- 5 1. A pharmaceutical tablet which comprises simvastatin and excipients, wherein the amount of lactose, if any, is less than 75 percent of the total of excipients by weight, and wherein the amount of cellulose is more than 20 percent of the total excipients by weight.
- 10 2. A tablet of claim 1 wherein the amount of lactose is less than 60 percent of the total of excipients by weight.
3. A tablet of claim 1 wherein the amount of lactose is less than 40 percent of the total of excipients by weight.
- 15 4. A tablet of claim 1 that is lactose free.
5. A tablet of any of claims 1 to 4 wherein the amount of cellulose is more than 40 percent of the total of excipients by weight.
- 20 6. A tablet of any of claims 1 to 4 wherein the amount of cellulose is more than 60 percent of the total of excipients by weight.
7. A tablet of any of claims 1 to 6 that comprises as a disintegrant other than cellulose.
- 25 8. A tablet of claim 7 wherein the disintegrant is starch.
9. A tablet of claim 8 wherein the amount of starch exceeds 12 percent of the total of excipients by weight.
- 30 10. A tablet of claim 8 wherein the amount of starch exceeds 20 percent of the total of excipients by weight.

11. A tablet of claim 7 wherein the disintegrant is crospovidone.
12. A tablet of claim 7 wherein the disintegrant is sodium starch glycolate.
- 5 13. A tablet of claim 7 wherein the disintegrant is either croscarmellose sodium or carmellose calcium.
- 10 14. A tablet of any of claims 11 to 13 wherein the amount of the disintegrant exceeds 1 percent of the total of excipients by weight.
- 15 15. A tablet of any of claims 11 to 13 wherein the amount of the disintegrant exceeds 2 percent of the total of excipients by weight.
- 16 16. A tablet of any of claims 11 to 13 wherein the amount of the disintegrant exceeds 3 percent of the total of the excipients by weight.
- 20 17. A tablet of any of claims 1 to 16 wherein excipients selected from the group consisting of cellulose, starch, sodium starch glycolate, and crospovidone exceed 65 percent of the total excipients by weight.
- 25 18. A tablet of any of claims 1 to 16 wherein excipients selected from the group consisting of cellulose, starch, sodium starch glycolate, and crospovidone exceed 80 percent of the total excipients by weight.
- 30 19. A tablet of any of claims 1 to 16 wherein excipients selected from the group consisting of cellulose, starch, sodium starch glycolate, and crospovidone exceed 90 percent of the total excipients by weight.
20. A tablet of any of claims 1 to 16 wherein excipients selected from the group consisting of cellulose, starch, sodium starch glycolate, and crospovidone exceed 95 percent of the total excipients by weight.

21. A tablet of any of claims 1 to 20 that is free of magnesium stearate.
22. A tablet of any of claims 1 to 21 that comprises zinc stearate.
- 5 23. A tablet of any of claims 1 to 21 that comprises sodium stearyl fumarate.
24. A tablet of any of claims 1 to 23 that is free of citric acid.
- 10 25. A tablet of any of claims 1 to 24 that is free of ascorbic acid.
26. A tablet of any of claims 1 to 25 that is free of butylated hydroxyanisole.
- 15 27. A tablet of any of claims 1 to 26, when made by a dry-mix process.

20

25

30

PCT/CA 03/00479

According to International Patent Classification (IPC) or to both national classification and IPC

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 62230 A (BAYER) 30 August 2001 (2001-08-30) claims ----	1-27
A	WO 99 30692 A (ANDRX) 24 June 1999 (1999-06-24) claims ----	1-27
A	GB 2 329 334 A (RECKITT & COLMAN) 24 March 1999 (1999-03-24) claims ----	1-27
A	WO 96 19201 A (MERCK) 27 June 1996 (1996-06-27) claims ----	1-27
	----- -/--	

X Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- | Special categories of cited documents: | |
|---|---|
| "A" document defining the general state of the art which is not considered to be of particular relevance | "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| "E" earlier document but published on or after the international filing date | "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. |
| "O" document referring to an oral disclosure, use, exhibition or other means | "G" document member of the same patent family |
| "P" document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search

18 June 2003

Date of mailing of the international search report

27/06/2003

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer _____

Scarponi, U

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 03/00479

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 915 954 A (A.D.AYER ET AL.) 10 April 1990 (1990-04-10) claims -----	1-27

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 03/00479

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0162230	A	30-08-2001	DE 10008506 A1	13-09-2001
			AU 3547001 A	03-09-2001
			WO 0162230 A1	30-08-2001
			EP 1259227 A1	27-11-2002
			US 2003031720 A1	13-02-2003
WO 9930692	A	24-06-1999	US 5916595 A	29-06-1999
			AU 737962 B2	06-09-2001
			AU 1627599 A	05-07-1999
			CA 2315329 A1	24-06-1999
			EP 1035842 A1	20-09-2000
			JP 2002508311 T	19-03-2002
			WO 9930692 A1	24-06-1999
GB 2329334	A	24-03-1999	NONE	
WO 9619201	A	27-06-1996	US 5582838 A	10-12-1996
			AU 693313 B2	25-06-1998
			AU 4472696 A	10-07-1996
			CA 2206211 A1	27-06-1996
			CN 1171048 A ,B	21-01-1998
			CZ 9701895 A3	18-02-1998
			EP 0801560 A1	22-10-1997
			FI 972586 A	17-06-1997
			HU 77370 A2	30-03-1998
			JP 11509829 T	31-08-1999
			NO 972880 A	20-06-1997
			NZ 298994 A	28-10-1998
			PL 320792 A1	27-10-1997
			RU 2168330 C2	10-06-2001
			SK 80597 A3	04-02-1998
			WO 9619201 A1	27-06-1996
US 4915954	A	10-04-1990	US 4814181 A	21-03-1989
			AT 400296 B	27-11-1995
			AT 214488 A	15-04-1995
			AU 2152688 A	09-03-1989
			BE 1001184 A3	08-08-1989
			CA 1288016 A1	27-08-1991
			CH 676794 A5	15-03-1991
			DE 3829942 A1	16-03-1989
			DK 485588 A	04-03-1989
			ES 2008580 A6	16-07-1989
			FI 884056 A	04-03-1989
			FR 2620025 A1	10-03-1989
			GB 2209280 A ,B	10-05-1989
			IE 61845 B1	30-11-1994
			IT 1223798 B	29-09-1990
			JP 1096116 A	14-04-1989
			KR 9506216 B1	12-06-1995
			LU 87326 A1	08-03-1989
			NL 8802180 A	03-04-1989
			NO 883837 A ,B,	06-03-1989
			PT 88404 A ,B	31-07-1989
			SE 8803084 A	04-03-1989
			US 4915953 A	10-04-1990
			ZA 8806435 A	30-05-1989